

Health Concerns About Dairy

PhysiciansCommittee
for Responsible Medicine

Many Americans, including some vegetarians, still consume substantial amounts of dairy products. And government policies still promote these products, despite scientific evidence that questions their health benefits and indicates their potential health risks. Though dairy is marketed as an essential food for strong bones, there is more to the story. Some important things to consider include potential health problems like heart disease, certain cancers, digestive problems, and type 1 diabetes.

Bone Health

Calcium is an important mineral that helps to keep bones strong. Our bones are constantly remodeling, meaning the body takes small amounts of calcium from the bones and replaces it with new calcium. Therefore, it is essential to have enough of this mineral so the body doesn't decrease bone density in this remodeling process. Though calcium is necessary for ensuring bone health, the actual benefits of consuming calcium diminish after a certain point. Research suggests that getting more than about 600 milligrams per day—easily achieved without dairy products or calcium supplements—does not make bones stronger.¹

In fact, research shows that dairy products have little or no benefit for bones. A 2005 review published in the journal *Pediatrics* showed that drinking milk does not improve bone strength in children.² In a more recent study, researchers tracked the diets, exercise, and stress fracture rates of young girls for seven years and concluded that dairy products and calcium do not prevent stress fractures in adolescent girls.³

How can this be? First, healthy bones need more than just calcium. For example, vitamin K is important for bone health, but most dairy products contain very little. Luckily, certain green vegetables such as kale and broccoli are especially beneficial for bones because they contain both calcium and vitamin K.

Vitamin D is also necessary for bone health. Without enough vitamin D, only 10-15 percent of the calcium you consume is absorbed.⁴ Milk does not naturally contain vitamin D. Rather, it's added to milk in a process called fortification. The natural source of vitamin D is sunlight: Our skin makes vitamin D when exposed to sun. Few foods naturally contain vitamin D—and no dairy products naturally contain this vitamin. Therefore, fortified cereals, bread, orange juice, and soy or other plant-based

milks exist as options for providing vitamin D through the diet.⁵ Supplements are also available.

Five to 15 minutes of midday sun exposure to the arms and legs, or the hands, face, and arms, can be enough to meet many people's vitamin D needs.⁶ However, having darker skin, being older, living in the north, living in an urban area, and even going through a dark winter season can all make it hard to get enough vitamin D from the sun alone. Luckily, supplements are an easy way to get vitamin D. The U.S. government recommends that adults 19-50 years old get 600 international units (IU) per day and that adults 51 years and older get 800 IU per day.

Other ways to protect bones include eating less salt,^{7,8} eating more fruits and vegetables,^{8,9} and ensuring adequate calcium intake from plant foods such as kale, broccoli, other leafy green vegetables, and beans. You can also use calcium-fortified products such as breakfast cereals and juices. Soy milk and fortified orange juice provide about the same amount of calcium per serving as milk or other dairy products.¹⁰

Lastly, exercise is one of the most effective ways to increase bone density and decrease the risk of osteoporosis.^{11,12} Its benefits have been observed in studies of both children and adults.^{11,13,14}

Fat Content and Heart Disease

Dairy products—including cheese, ice cream, milk, butter, and yogurt—contribute significant amounts of cholesterol and saturated fat to the diet. Diets high in fat, and especially in saturated fat, can increase the risk of heart disease and stroke and can cause other serious health problems.^{15,16}

On the other hand, a low-fat, plant-based diet (which eliminates dairy products), in combination with exercise, smoking cessation, and stress management, can not only prevent heart disease, but may even reverse it.^{17,18}

Cancer

Consumption of dairy products has also been linked to higher risk for various cancers, especially to cancers of the reproductive system. Most significantly, consuming dairy has been linked to increased risk for prostate cancer.¹⁹⁻²¹

The danger of dairy product consumption as it relates to prostate

cancer is most likely linked to increases in insulin-like growth factor (IGF-1).²² Consuming milk and dairy products on a regular basis has been shown to increase blood levels of IGF-1 in humans.^{23,24} Studies in diverse populations have shown a strong and consistent link between IGF-1 in the blood and prostate cancer risk.^{25,26} One study showed that men with the highest levels of IGF-1 had more than four times the risk of prostate cancer, compared with those who had the lowest levels.²⁷ In the Physicians Health Study, which tracked 21,660 participants for 28 years, researchers found an increased risk of prostate cancer for those who consumed more than 2.5 servings of dairy products per day as compared with those who had fewer than 0.5 servings

Certain **green vegetables** such as kale and broccoli are especially beneficial for bones because they **contain both calcium and vitamin K.**

a day.²⁰ This study, which is supported by other findings,^{21,28,29} also shows that prostate cancer risk is higher with increased consumption of low-fat milk. That means that too much dairy calcium, and not just the fat in dairy products, could harm prostate health.

A study of 1,893 women diagnosed with early-stage invasive breast cancer revealed that eating more high-fat dairy products was linked to higher mortality (death) rates. As little as half a serving per day increased risk significantly. Since hormones are stored in fat, consuming high-fat, rather than low-fat, dairy products likely means women are consuming more estrogen.³⁰ A second large study of 1,941 women found that women who consumed the highest amounts of cheddar, American, and cream cheeses had a 53 percent higher risk for breast cancer.



The consumption of dairy products may also contribute to development of ovarian cancer. The relationship between dairy products and ovarian cancer may be due to the breakdown of the milk sugar (lactose) into galactose, a sugar which may be toxic to ovarian cells.³¹ Two studies, one conducted in Sweden and one conducted among African-American women, showed that consuming lactose and dairy products was positively linked to ovarian cancer.^{32,33} The Iowa Women's Health Study, found that women who consumed more than one glass of milk per day had a 73 percent greater chance of developing ovarian cancer than women who drank less than one glass per day.³⁴

Lastly, a large study published in the *British Journal of Cancer* identified 22,788 people who were lactose intolerant and found that those who avoided dairy (due to lactose intolerance) had a lower incidence of lung, breast, and ovarian cancers than their family members who did not avoid dairy. The researchers suggest that avoiding the saturated fat and extra hormones found in dairy products is protective against certain types of cancer.³⁵

Lactose Intolerance

Lactose intolerance is common, affecting about 95 percent of Asian Americans, 74 percent of Native Americans, 70 percent of African Americans, 53 percent of Mexican Americans, and 15 percent of Caucasians.³⁶ Symptoms, which include upset stomach, diarrhea, and gas, occur because these individuals lack the enzyme lactase, which is needed to digest the milk sugar lactose. Nursing children make enzymes that break down lactose, but as we grow up, many of us lose this capacity.³⁷ As a result, lactose is not absorbed, but remains in the intestine where it causes symptoms.

Contaminants

Dairy contains contaminants that range from hormones to pesticides. Milk naturally contains hormones and growth factors produced within a cow's body, too. In addition, man-made hormones such as recombinant bovine growth hormone (rBGH) are commonly given to cows to increase their milk production.³⁸

Antibiotics are used to treat udder infections (mastitis) in cows. Traces of these antibiotics have been found in some samples of milk and dairy products.³⁹ Unfortunately, antibiotics are used frequently, because mastitis is very common in cows due to farming practices that cause cows to produce more milk than nature intended.

Pesticides, polychlorinated biphenyls (PCBs), and dioxins are other examples of contaminants found in milk. Dairy products contribute to one-fourth to one-half of the dietary intake of total dioxins.⁴⁰ All these toxins tend to build up in the body over time. Eventually, this can harm the immune, reproductive, and nervous systems. Moreover, PCBs and dioxins have been linked to cancer.⁴¹

Other contaminants that can make their way into milk include melamine (often found in plastics and can harm the kidneys and urinary tract) and cancer-causing toxins like aflatoxins.^{42,43}

Milk Proteins and Diabetes

Insulin-dependent (type 1) diabetes is linked to consumption of dairy products in infancy.⁴⁴ A 2001 Finnish study of nearly 3,000 infants with genetically increased risk for developing diabetes showed that early introduction of cow's milk increased susceptibility to type 1 diabetes.⁴⁵ In addition, the American Academy of Pediatrics observed up to a 30 percent reduction in the incidence of type 1 diabetes in infants who avoid exposure to cow's milk protein for at least the first three months of their lives.⁴⁶

Health Concerns for Children and Infants

Milk proteins, milk sugar, fat, and saturated fat in dairy products pose health risks for children and can lead to obesity, diabetes, and heart disease. While low-fat milk is often recommended for decreasing obesity risk, a study published in the *Archives of Disease in Childhood* showed that children who drank 1 percent or skim milk, compared with those who drank full-fat milk, were not any less likely to be obese.⁴⁷ Moreover, a meta-analysis found no support for the argument that increasing dairy product intake decreases body fat and weight over the long term (>1 year).⁴⁸

The consumption of cow's milk is not recommended for infants. The American Academy of Pediatrics recommends that infants

below one year of age not be given cow's milk.⁴⁹ Not only is the iron in cow's milk hard for babies to absorb, but whole cow's milk can cause microscopic bleeding in infants' intestines.⁵⁰

Colic is an additional concern with milk consumption. Up to 28 percent of infants suffer from colic during the first month of life.⁵¹ Pediatricians learned long ago that cow's milk was often the reason. We now know that breastfeeding mothers can have colicky babies if the *mothers* consume cow's milk. Cow's milk proteins can pass through the mother's bloodstream, into her breast milk, and to the baby, causing symptoms in some infants.^{52,53}

Additionally, children who consume cow's milk are more likely to develop food allergies^{54,55} and are more likely to suffer from chronic constipation.^{56,57}

Conclusions

Milk and dairy products are not necessary in the diet and can even be harmful to health. It's best to consume a healthful diet of whole grains, fruits, vegetables, beans, peas, and lentils and replace cow's milk with nondairy milks like almond, soy, or cashew milk. These nutrient-dense foods can help you meet your nutrient requirements with ease—and without the health risks associated with dairy products. ◀

References

- Feskanih D, Willett WC, Colditz GA. Calcium, vitamin D, milk consumption, and hip fractures: a prospective study among postmenopausal women. *Am J Clin Nutr.* 2003;77:504–511.
- Lanou AJ, Berkow SE, Barnard ND. Calcium, dairy products, and bone health in children and young adults: a reevaluation of the evidence. *Pediatrics.* 2005;115:736–743.
- Sonneville KR, Gordon CM, Kocher MS, Pierce LM, Ramappa A, Field AE. Vitamin D, calcium, and dairy intakes and stress fractures among female adolescents. *Arch Pediatr Adolesc Med.* 2012;166:595–600.
- Holick MF, Garabedian M. Vitamin D: photobiology, metabolism, mechanism of action, and clinical applications. In: Favus MJ, ed. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism.* 6th ed. Washington, D.C.: American Society for Bone and Mineral Research; 2006:129–137.
- Zhang R, Naughton D. Vitamin D in health and disease: current perspectives. *Nutr J.* 2010;9:65–78.
- Holick M. The vitamin D epidemic and its health consequences. *J Nutr.* 2005;135:2739S–2748S.
- Reid DM, New SA. Nutritional influences on bone mass. *Proceed Nutr Soc.* 1997;56:977–987.
- Lin P, Ginty F, Appel L, et al. The DASH diet and sodium reduction improve markers of bone turnover and calcium metabolism in adults. *J Nutr.* 2001;133:3130–3136.
- Qiu R, Cao W, Tian H, He J, Chen G, Chen Y. Greater intake of fruit and vegetables is associated with greater bone mineral density and lower osteoporosis risk in middle-aged and elderly adults. *PLoS ONE.* 2017;12:e0168906–e0168919.
- National Institutes of Health. NIH Osteoporosis and Related Bone Diseases National Resources Center. Calcium and Vitamin D: Important at Every Age. <https://www.bones.nih.gov/health-info/bone/bone-health/nutrition/calcium-and-vitamin-d-important-every-age>. Accessed May 20, 2018.
- Marques EA, Mota J, Carvalho J. Exercise effects on bone mineral density in older adults: a meta-analysis of randomized controlled trials. *Age.* 2012;34:1493–1515.
- Going S, Lohman T, Houtkooper L, et al. Effects of exercise on bone mineral density in calcium-replete postmenopausal women with and without hormone replacement therapy. *Osteoporos Int.* 2003;14:637–643.
- Lunt M, Masaryk P, Scheidt-Nave C, et al. The effects of lifestyle, dietary dairy intake and diabetes on bone density and vertebral deformity prevalence: the EVOS study. *Osteoporos Int.* 2001;12:688–698.
- Lloyd T, Beck TJ, Lin HM, et al. Modifiable determinants of bone status in young women. *Bone.* 2002;30:416–421.
- Li Y, Hruby A, Bernstein AM, et al. Saturated fats compared with unsaturated fats and sources of carbohydrates in relation to risk of coronary heart disease: a prospective cohort study. *J Am Coll Cardiol.* 2015;66:1538–1548.
- Chen M, Li Y, Sun Q, et al. Dairy fat and risk of cardiovascular disease in 3 cohorts of US adults. *Am J Clin Nutr.* 2016;104:1209–1217.
- Szeto YT, Kwok TC, Benzie IF. Effects of a long-term vegetarian diet on biomarkers of antioxidants status and cardiovascular disease risk. *Nutrition.* 2004;20:863–866.
- Ornish D, Brown SE, Scherwitz LW, et al. Can lifestyle changes reverse coronary heart disease? *Lancet.* 1990;336:129–133.
- Qin L, Xu J, Wang P, Tong J, Hoshi K. Milk consumption is a risk factor for prostate cancer in Western countries: evidence from cohort studies. *Asia Pac J Clin Nutr.* 2007;16:467–476.
- Song Y, Chavarro JE, Cao Y, et al. Whole milk intake is associated with prostate cancer-specific mortality among U.S. male physicians. *J Nutr.* 2013;143:189–196.
- Chan JM, Stampfer MJ, Ma J, Gann PH, Gaziano JM, Giovannucci E. Dairy products, calcium, and prostate cancer risk in the Physicians' Health Study. *Am J Clin Nutr.* 2001;74:549–554.

22. Voskuil DW, Vrieling A, van't Veer LJ, Kampman E, Rookus MA. The insulin-like growth factor system in cancer prevention: potential of dietary intervention strategies. *Cancer Epidemiol Biomarkers Prev.* 2005;14:195–203.
23. Qin LQ, He K, Xu JY. Milk consumption and circulating insulin-like growth factor-I level: a systematic literature review. *Int J Food Sci Nutr.* 2009;60 Suppl 7:330–340.
24. Qin LQ, He K, Xu JY. Milk consumption and circulating insulin-like growth factor-I level: a systematic literature review. *Int J Food Sci Nutr.* 2009;60:330–340.
25. Cohen P. Serum insulin-like growth factor-I levels and prostate cancer risk—interpreting the evidence. *J Natl Cancer Inst.* 1998;90:876–879.
26. Shi R, Berkel HJ, Yu H. Insulin-like growth factor-I and prostate cancer: a meta-analysis. *Br J Cancer.* 2001;85:991–996.
27. Chan JM, Stampfer MJ, Giovannucci E, et al. Plasma insulin-like growth factor-1 and prostate cancer risk: a prospective study. *Science.* 1998;279:563–565.
28. Tseng M, Breslow RA, Graubard BI, Ziegler RG. Dairy, calcium and vitamin D intakes and prostate cancer risk in the National Health and Nutrition Examination Epidemiologic Follow-up Study cohort. *Am J Clin Nutr.* 2005;81:1147–1154.
29. Yang M, Kenfield SA, Van Blarigan EL, et al. Dairy intake after prostate cancer diagnosis in relation to disease-specific and total mortality. *Int J Cancer.* 2015;137:2462–2469.
30. Kroenke CH, Kwan ML, Sweeney C, Castillo A, Caan BJ. High- and low-fat dairy intake, recurrence, and mortality after breast cancer diagnosis. *J Natl Cancer Inst.* 2013;105:616–623.
31. Cramer DW, Greenberg ER, Titus-Ernstoff L, et al. A case-control study of galactose consumption and metabolism in relation to ovarian cancer. *Cancer Epidemiol Biomarkers Prev.* 2000;9:95–101.
32. Larsson SC, Bergkvist L, Wolk A. Milk and lactose intakes and ovarian cancer risk in the Swedish Mammography Cohort. *Am J Clin Nutr.* 2004;80:1353–1357.
33. Qin B, Moorman PG, Alberg AJ, et al. Dairy, calcium, vitamin D and ovarian cancer risk in African American Women. *Br J Cancer.* 2016;115:1122–1130.
34. Kushi LH, Mink PJ, Folsom AR, et al. Prospective study of diet and ovarian cancer. *Am J Epidemiol.* 1999;149:21–31.
35. Ji J, Sundquist J, Sundquist K. Lactose intolerance and risk of lung, breast and ovarian cancers: aetiological clues from a population-based study in Sweden. *Br J Cancer.* 2015;112:149–152.
36. Bertron P, Barnard ND, Mills M. Racial bias in federal nutrition policy, part I: the public health implications of variations in lactase persistence. *J Natl Med Assoc.* 1999;91:151–157.
37. Swallow DM. Genetics of lactase persistence and lactose intolerance. *Annu Rev Genet.* 2003;37:197–219.
38. Outwater JL, Nicholson A, Barnard N. Dairy products and breast cancer: the IGF-1, estrogen, and bGH hypothesis. *Med Hypothesis.* 1997;48:453–461.
39. Azzouz A, Jurado-Sánchez B, Souhail B, Ballesteros E. Simultaneous determination of 20 pharmacologically active substances in cow's milk, goat's milk, and human breast milk by gas chromatography–mass spectrometry. *J Agric Food Chem.* 2011;59:5125–5132.
40. Bhandari SD, Schmidt RH, Rodrick GE. Hazards resulting from environmental, industrial, and agricultural contaminants. In: Schmidt RH, Rodrick GE, eds. *Food Safety Handbook.* Hoboken, NJ: John Wiley & Sons, Inc.; 2005:291–321.
41. National Institutes of Environmental Health Sciences. Health and Education: Dioxins. Available at: <https://www.niehs.nih.gov/health/topics/agents/dioxins/index.cfm>. Reviewed August 28, 2017. Accessed May 21, 2018.
42. Fischer WJ, Schilter B, Tritscher AM, Stadler RH. Contaminants of milk and dairy products: contamination resulting from farm and dairy practices. In: Fuquay JW, ed. *Encyclopedia of Dairy Sciences.* 2nd ed. San Diego, CA: Academic Press; 2011:887–897.
43. Prandini A, Tansini G, Sigolo S, Filippi L, Laporta M, Piva G. On the occurrence of aflatoxin M1 in milk and dairy products. *Food Chem Toxicol.* 2009;47:984–991.
44. Chia JSJ, McRae JL, Kukuljan S, et al. A1 beta-casein milk protein and other environmental pre-disposing factors for type 1 diabetes. *Nutr Diabetes.* 2017;7:e274–e281.
45. Kimpimaki T, Erkkola M, Korhonen S, et al. Short-term exclusive breastfeeding predisposes young children with increased genetic risk of type 1 diabetes to progressive beta-cell autoimmunity. *Diabetologia.* 2001;44:63–69.
46. Eidelman AI, Schanler RJ. Policy statement: breastfeeding and the use of human milk. From the American Academy of Pediatrics. *Pediatrics.* 2012;129:827–841.
47. Scharf RJ, Demmer RT, DeBoer MD. Longitudinal evaluation of milk type consumed and weight status in preschoolers. *Arch Dis Child.* 2013;98:335–340.
48. Chen M, Pan A, Malik VS, Hu FB. Effects of dairy intake on body weight and fat: a meta-analysis of randomized controlled trials. *Am J Clin Nutr.* 2012;96:735–747.
49. Gartner LM, Morton J, Lawrence RA, et al. Breastfeeding and the use of human milk. *Pediatrics.* 2005;115:496–506.
50. Leung AK, Sauve RS. Whole cow's milk in infancy. *Paediatr Child Health.* 2003;8:419–421.
51. Lucassen PL, Assendelft WJ, van Eijk JT, Gubbels JW, Douwes AC, van Geldrop WJ. Systematic review of the occurrence of infantile colic in the community. *Arch Dis Child.* 2001;84:398–403.
52. Jarvinen KM, Makinen-Kiljunen S, Suomalainen H. Cow's milk challenge through human milk evoked immune responses in infants with cow's milk allergy. *J Pediatr.* 1999;135:506–512.
53. Leung AKC. Dietary manipulations for infantile colic. *Paediatr Child Health.* 2003;8:449–452.
54. Sampson HA. Update on food allergy. *J Allergy Clin Immunol.* 2004;113:805–819.
55. Host A. Frequency of cow's milk allergy in childhood. *Ann Allergy Asthma Immunol.* 2002;89(6 Suppl 1):33–37.
56. Iacono G, Cavataio F, Montalto G, et al. Intolerance of cow's milk and chronic constipation in children. *N Engl J Med.* 1998;339:1100–1104.
57. Dehghani SM, Ahmadvour B, Haghghat M, Kashef S, Imanieh MH, Soleimani M. The role of cow's milk allergy in pediatric chronic constipation: a randomized clinical trial. *Iran J Pediatr.* 2012;22:468–474.

